arXiv:1302.7118v2 [math.CV] 18 Mar 2013

Asymptotics of the Discrete Chebyshev Polynomials

J. H. Pan[a] and R. Wong[b]

[a] *Department of Mathematics, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong*

[b] *Liu Bie Ju Centre for Mathematical Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong*

Abstract

The discrete Chebyshev polynomials $t_n(x, N)$ are orthogonal with respect to a distribution, which is a step function with jumps one unit at the points $x = 0, 1, \cdots, N-1$, N being a fixed positive integer. By using a double integral representation, we have recently obtained asymptotic expansions for $t_n(aN, N+1)$ in the double scaling limit, namely, $N \to \infty$ and $n/N \to b$, where $b \in (0, 1)$ and $a \in (-\infty, \infty)$; see [Studies in Appl. Math. **128** (2012), 337-384]. In the present paper, we continue to investigate the behaviour of these polynomials when the parameter b approaches the endpoints of the interval $(0, 1)$. While the case $b \to 1$ is relatively simple (since it is very much like the case when b is fixed), the case $b \to 0$ is quite complicated. The discussion of the latter case is divided into several subcases, depending on the quantities n, x and xN/n^2, and different special functions have been used as approximants, including Airy, Bessel and Kummer functions.

1 INTRODUCTION

The discrete Chebyshev polynomials $t_n(x, N)$ can be defined as a special case of the Hahn polynomials [1, p.174]

$$\begin{aligned} Q_n(x;\alpha,\beta,N) &= {}_3F_2(-n,-x,n+\alpha+\beta+1;-N,\alpha+1;1) \\ &= \sum_{k=0}^{n} \frac{(-n)_k(-x)_k(n+\alpha+\beta+1)_k}{(-N)_k(\alpha+1)_k k!}. \end{aligned} \tag{1.1}$$

With $\alpha = \beta = 0$ in (1.1), we have

$$\begin{aligned} t_n(x,N) &= (-1)^n(N-n)_n Q_n(x;0,0,N-1) \\ &= (-1)^n(N-n)_n \sum_{k=0}^{n} \frac{(-n)_k(-x)_k(n+1)_k}{(-N+1)_k k!k!}; \end{aligned} \tag{1.2}$$

see [1, p.176].

In a recent paper [8], we have studied the asymptotic behaviour of $t_n(x, N+1)$ as n, $N \to \infty$ in such a way that the ratios

$$a = x/N \qquad \text{and} \qquad b = n/N, \tag{1.3}$$

satisfy the inequalities

$$-\infty < a \le \frac{1}{2} \qquad \text{and} \qquad 0 < b < 1. \tag{1.4}$$

In view of the symmetry relation [8, eq.(8)]

$$t_n(x, N+1) = (-1)^n t_n(N-x, N+1), \tag{1.5}$$

our study actually covers the entire real-axis $-\infty < x < \infty$ or, equivalently, the entire parameter range $-\infty < a < \infty$. For a discussion of the asymptotic behaviour of the Hahn polynomials, see [4].

In the present paper, we shall investigate the behaviour of the polynomials $t_n(x, N+1)$, when the parameter b given in (1.3) either tends to 0 or tends to 1. The case $b \to 1$ turns out to be rather simple, since the ultimate expansions and their derivations are similar to those in the case when b is fixed; see (2.21) for $0 \le a \le \frac{1}{2}$ and (2.28) for $a < 0$.

To derive asymptotic approximation for $t_n(x, N+1)$ when $b \to 0$, we divide our discussion into several cases, depending on the quantities n, x and xN/n^2. A summary of our findings is given in Table 1 below. Some explanation is needed for this table. By "small", "fixed" and "large", we mean, respectively, $x = o(1)$, $x \in [\delta, M]$ and $x \gg O(1)$ as $N \to \infty$, where δ is a small positive number and M is a large positive number. By "Airy", we mean an asymptotic expansion whose associated approximants are the Airy functions $\mathrm{Ai}(z)$ and $\mathrm{Bi}(z)$. In the same manner, by "Bessel" and "Kummer", we mean

Table 1: Asymptotic expansions in different cases when $\frac{n}{N} \to 0$ and $N \to \infty$

x \ xN/n^2	small	fixed	large
large	Kummer	Airy	Bessel
fixed	Kummer	Airy	series(1.2)
small	Kummer / series(1.2)	Airy	series(1.2)

asymptotic expansions whose associated approximants are, respectively, the Bessel function $J_v(z)$ and the Kummer function $M(a,c,z)$. The Kummer function used in this paper is defined by

$$\mathbf{M}(a,c,z) = \frac{1}{\Gamma(c)} M(a,c,z) = \frac{1}{\Gamma(c)} \sum_{s=0}^{\infty} \frac{(a)_s}{(c)_s} \frac{z^s}{s!}, \tag{1.6}$$

which was introduced in [6, p.255]. The case involving Bessel functions covers an earlier result of Sharapodinov [10], where, instead of Bessel functions, Jacobi polynomials are used as an approximant. In the case when xN/n^2 is large and x is either fixed or small, the series in (1.2) is itself an asymptotic expansion (in the generalized sense [11, p.10]) as $N \to \infty$. Hence, there is no need to seek for another asymptotic representation.

The arrangement of the present paper is as follows. In Section 2, we recall the major results in [8]; to facilitate our presentation for later sections, we also include here brief derivations of the expansions given in [8]. In Section 3, we consider the case when xN/n^2 is small and show that the expansions are of Kummer-type. In Section 4, we consider the case when the quantity xN/n^2 is fixed; that is, when xN/n^2 is bounded away from zero and infinity. There are three subcases, depending on x being large, fixed or small. In all three subcases, we will show that the expansions are of Airy-type; see Table 1. The case when both quantities x and xN/n^2 are large is dealt with in Section 5, where we show that the expansion of $t_n(x, N+1)$ can be expressed in terms of Bessel functions. The final section is devoted to two remaining cases, namely, (i) when xN/n^2 is large and x is either fixed or small, and (ii) $x < 0$.

2 RESULTS IN REFERENCE [8]

In [8], we have divided our discussion into two cases: (i) $0 \le a \le \frac{1}{2}$, and (ii) $a < 0$. As mentioned in Section 1, by virtue of the symmetry relation (1.5), these two cases cover the entire range $-\infty < a < \infty$.

In case (i), we started with the integral representation

$$t_n(x, N+1) = \frac{(-1)^n}{2\pi i} \frac{\Gamma(n+N+2)}{\Gamma(n+1)\Gamma(N-n+1)} \int_0^1 \int_{\gamma_1} \frac{1}{w-1} e^{Nf(t,w)} \mathrm{d}w \mathrm{d}t, \tag{2.1}$$

where

$$f(t,w) = b\ln(1-t) + (1-b)\ln t + a\ln w - a\ln(w-1) + b\ln[1-(1-t)w], \tag{2.2}$$

and the curve γ_1 starts at $w = 0$, runs along the lower edge of the positive real line towards $w = 1$, encircles the point $w = 1$ in the counterclockwise direction and returns to the origin along the upper edge of the positive real line. As a function of two variables, the partial derivatives $\partial f/\partial t$ and $\partial f/\partial w$ vanish at $(t, w) = (t_\pm, w_\pm)$, where

$$t_+ = \frac{2 - 2a - b^2 + b\sqrt{b^2 - 4a + 4a^2}}{2(1-a)(1+b)}, \qquad w_+ = \frac{b + \sqrt{b^2 - 4a + 4a^2}}{2b}, \tag{2.3}$$

and

$$t_- = \frac{2 - 2a - b^2 - b\sqrt{b^2 - 4a + 4a^2}}{2(1-a)(1+b)}, \qquad w_- = \frac{b - \sqrt{b^2 - 4a + 4a^2}}{2b}. \tag{2.4}$$

For each fixed $w \in \gamma_1$, we now find a steepest descent path of the phase function $f(t, w)$ in the variable t, which passes through a saddle point $t_0(w)$ depending on w. (Note that not only the saddle point $t_0(w)$, but all points t on this steepest descent path depend on w.) The function $f(t_0(w), w)$ is a function of w alone. To find the relevant saddle point $t_0(w)$, we solve the equation $\partial f(t, w)/\partial t = 0$, and obtain

$$t_0(w) = \frac{2w - 1 + \sqrt{1 + 4b^2(w-1)w}}{2(1+b)w}. \tag{2.5}$$

It can be shown that

$$t_0(w_+) = t_+, \qquad t_0(w_-) = t_-. \tag{2.6}$$

Define the standard transformation $t \to \tau$ by

$$\tau^2 = f(t_0(w), w) - f(t, w); \tag{2.7}$$

see [11, p.88]. Note that we have $\tau = -\infty$ when $t = 0$, and $\tau = +\infty$ when $t = 1$. Furthermore, this mapping takes $t = t_0(w)$ to $\tau = 0$. Coupling (2.1) and (2.7) gives

$$\begin{aligned} t_n(x, N+1) =& \frac{(-1)^n}{2\pi i} \frac{\Gamma(n+N+2)}{\Gamma(n+1)\Gamma(N-n+1)} \\ & \times \int_{-\infty}^{+\infty} \int_{\gamma_1} \frac{1}{w-1} e^{Nf(t_0(w),w)} e^{-N\tau^2} \frac{\mathrm{d}t}{\mathrm{d}\tau} \mathrm{d}w \mathrm{d}\tau, \end{aligned} \tag{2.8}$$

where γ_1 is the integration path of the variable w in (2.1). Note that the first variable of f in (2.8) is now $t_0(w)$, instead of t. Hence, the phase function is a function of w alone. Setting $\partial f(t_0(w), w)/\partial w = 0$, we obtain the saddle points

$$w_\pm = \frac{b \pm \sqrt{b^2 - 4a + 4a^2}}{2b}; \tag{2.9}$$

cf.(2.3) and (2.4). Motivated by (2.2), define the new phase function

$$\psi(u) = a \ln u - a \ln(u-1) + \eta u. \tag{2.10}$$

The saddle points of ψ are given by

$$u_\pm = \frac{\eta \pm \sqrt{\eta^2 + 4a\eta}}{2\eta}. \tag{2.11}$$

To reduce the double integral in (2.1) into a canonical form, we define the second mapping $w \to u$ by

$$\begin{aligned} f(t_0(w), w) &= \psi(u) + \gamma \\ &= a \ln u - a \ln(u-1) + \eta u + \gamma \end{aligned} \tag{2.12}$$

with

$$u(w_+) = u_-, \qquad u(w_-) = u_+, \tag{2.13}$$

where η and γ are real numbers depending on the parameters a and b in (2.2). From (2.12), we have

$$\begin{aligned} \frac{\mathrm{d}w}{\mathrm{d}u} &= \frac{\mathrm{d}\psi}{\mathrm{d}u} \Big/ \frac{\mathrm{d}f(t_0(w), w)}{\mathrm{d}w} \\ &= \frac{\eta(u-u_+)(u-u_-)w(1-w)\left[(2a-1) - \sqrt{1+4b^2w^2-4b^2w}\right]}{-2b^2u(u-1)(w-w_+)(w-w_-)} \end{aligned} \tag{2.14}$$

for $u \neq u_\pm$, and by L'Hôpital's rule,

$$\frac{\mathrm{d}w}{\mathrm{d}u} = \left[\frac{\sqrt{\eta^2+4a\eta}(1-2a)(1-a)(-\eta)}{b^3\sqrt{b^2-4a+4a^2}}\right]^{1/2} \tag{2.15}$$

for $u = u_\pm$. With the change of variable $w \to u$ defined in (2.12), the representation in (2.8) becomes

$$\begin{aligned} t_n(x, N+1) =& \frac{(-1)^n}{2\pi i} \frac{\Gamma(n+N+2)}{\Gamma(n+1)\Gamma(N-n+1)} e^{N\gamma} \\ &\times \int_{-\infty}^{+\infty} \int_{\gamma_u} \frac{h(u,\tau)}{u-1} e^{N\psi(u)-N\tau^2} \mathrm{d}u\mathrm{d}\tau, \end{aligned} \tag{2.16}$$

where

$$h(u,\tau) = \frac{u-1}{w-1} \frac{\mathrm{d}w}{\mathrm{d}u} \frac{\mathrm{d}t}{\mathrm{d}\tau}, \tag{2.17}$$

and γ_u is the steepest descent path of $\psi(u)$ in the u-plane. An asymptotic expansion, holding uniformly for $a \in [0, \frac{1}{2}]$, was then derived in [8] by an integration-by-parts technique. To state the result, we define recursively $h_0(u,\tau) = h(u,\tau)$,

$$h_l(u,\tau) = a_l(\tau) + b_l(\tau)u - (u-u_-)(u-u_+)g_l(u,\tau) \tag{2.18}$$

and

$$h_{l+1}(u,\tau) = (u-1)\left[g_l(u,\tau) + u\frac{\partial g_l(u,\tau)}{\partial u}\right], \qquad l = 0, 1, 2, ... \tag{2.19}$$

Furthermore, we write

$$a_l(\tau) = \sum_{j=0}^{\infty} a_{l,j}\tau^j, \qquad b_l(\tau) = \sum_{j=0}^{\infty} b_{l,j}\tau^j. \tag{2.20}$$

The resulting expansion takes the form

$$t_n(x,N+1)\sim\frac{(-1)^n\Gamma(n+N+2)e^{N\gamma}}{\Gamma(n+1)\Gamma(N-n+1)\sqrt{N}}\left\{\mathbf{M}(aN+1,1,\eta N)\sum_{l=0}^{\infty}\frac{c_l}{(\eta N)^l}\right.$$
$$\left.+\,\mathbf{M}'(aN+1,1,\eta N)\sum_{l=0}^{\infty}\frac{d_l}{(\eta N)^l}\right\},\tag{2.21}$$

where $\mathbf{M}(a,c,z)$ is the Kummer function defined in (1.6), and the coefficients c_l and d_l are given explicitly by

$$c_l=\sum_{m=0}^{l}a_{l-m,2m}\Gamma(m+\frac{1}{2})\eta^m,\qquad d_l=\sum_{m=0}^{l}b_{l-m,2m}\Gamma(m+\frac{1}{2})\eta^m.\tag{2.22}$$

Case (ii) was dealt with in a similar manner. We started with the double-integral representation

$$t_n(x,N+1)=\frac{(-1)^n}{2\pi i}\frac{\Gamma(n+N+2)}{\Gamma(n+1)\Gamma(N-n+1)}\int_0^1\int_{\gamma_2}\frac{1}{w-1}e^{N\widetilde{f}(t,w)}\mathrm{d}w\mathrm{d}t,\tag{2.23}$$

where

$$\widetilde{f}(t,w)=b\ln(1-t)+(1-b)\ln t+a\ln(-w)-a\ln(1-w)+b\ln[1-(1-t)w],\tag{2.24}$$

and the integration path γ_2 starts at $w=1$, traverses along the upper edge of the positive real line towards $w=0$, encircles the origin in the counterclockwise direction and returns to $w=1$ along the lower edge of the positive real line.

Now we follow the same argument as given in Case (i), and use the same mapping $t\to\tau$ defined in (2.7), except with $f(t,w)$ replaced by $\widetilde{f}(t,w)$. The result is

$$t_n(x,N+1)=\frac{(-1)^n}{2\pi i}\frac{\Gamma(n+N+2)}{\Gamma(n+1)\Gamma(N-n+1)}$$
$$\times\int_{-\infty}^{+\infty}\int_{\gamma_2}\frac{1}{w-1}e^{N\widetilde{f}(t_0(w),w)}e^{-N\tau^2}\frac{\mathrm{d}t}{\mathrm{d}\tau}\mathrm{d}w\mathrm{d}\tau.\tag{2.25}$$

Because of the shape of the contour γ_2, it turns out that the phase function $\widetilde{f}(t_0(w),w)$ has only one relevant saddle point, namely w_-; see (2.9). As a consequence, we define the mapping $w\to u$ by

$$\widetilde{f}(t_0(w),w)=a\ln(-u)-u+\gamma\tag{2.26}$$

with

$$u(w_-)=a,\tag{2.27}$$

where γ is a constant depending on the parameters a and b in (1.3). The final expansion is in terms of the gamma function, and we have

$$t_n(x,N+1)\sim\frac{(-1)^n\Gamma(n+N+2)N^{-aN}e^{N\gamma}}{\Gamma(n+1)\Gamma(N-n+1)\Gamma(-aN+1)}\sum_{l=0}^{\infty}\frac{c_l}{N^{l+\frac{1}{2}}},\tag{2.28}$$

where c_l are constants that can be given recursively.

3 KUMMER-TYPE EXPANSION

As $b \to 0$, some of the steps in Section 2 are no longer valid. Let us first examine the mapping $t \to \tau$ given in (2.7). In this mapping, we have used the fact that for each fixed $w \in \gamma_1$, the saddle point $t = t_0(w)$ in (2.5) is bounded away from $t = 0$ and $t = 1$, and the steepest descent path from $t = 0$ to $t = 1$, passing through $t_0(w)$, is mapped onto $(-\infty, \infty)$ in the τ-plane. However, from (2.5), we have for any fixed w

$$t_0(w) = 1 - b + O\left(b^2\right) \qquad \text{as } b \to 0; \tag{3.1}$$

that is, as $b \to 0$, $t_0(w)$ approaches the branch point $t = 1$ in the t-plane. Thus, the mapping (2.7) is no longer suitable in this case. Since we are more interested in the neighbourhood of $t = 1$, where the term $b\ln(1-t)$ in the phase function (2.2) becomes singular, we introduce the mapping

$$f(t, w) = b\ln\tau - \tau + A, \tag{3.2}$$

where the constant A does not depend on t or τ (but may depend on w); see (3.4) below. To make the mapping $t \to \tau$ defined in (3.2) one-to-one and analytic, we prescribe $t = t_0(w)$ to correspond to $\tau = b$, which is the saddle point of $b\ln\tau - \tau + A$; i.e.,

$$\tau(t_0(w)) = b. \tag{3.3}$$

This gives

$$A = f(t_0(w), w) - b\ln b + b. \tag{3.4}$$

Note that we have $\tau = 0$ when $t = 1$, and $\tau = +\infty$ when $t = 0$. Furthermore, from (3.2), we have

$$\begin{aligned} \frac{\mathrm{d}t}{\mathrm{d}\tau} &= \frac{b/\tau - 1}{\partial f(t,w)/\partial t} = \frac{(\tau - b)t(1-t)[1-(1-t)w]}{\tau(1+b)w(t - t_0^+(w))(t - t_0^-(w))}, && \tau \neq b, \\ \frac{\mathrm{d}t}{\mathrm{d}\tau} &= \left\{ \frac{t_0^+(w)(1-t_0^+(w))[1-(1-t_0^+(w))w]}{b\sqrt{1+4b^2(w-1)w}} \right\}^{1/2}, && \tau = b, \end{aligned} \tag{3.5}$$

where we have used L'Hôpital's rule for $\tau = b$ and

$$t_0^{\pm}(w) = \frac{2w - 1 \pm \sqrt{1+4b^2(w-1)w}}{2(1+b)w};$$

cf.(2.5). Coupling (2.1) and (3.2), we have, instead of (2.8),

$$\begin{aligned} t_n(x, N+1) =& \frac{(-1)^{n+1}}{2\pi i} \frac{\Gamma(n+N+2)e^n b^{-n}}{\Gamma(n+1)\Gamma(N-n+1)} \\ & \times \int_0^{+\infty} \int_{\gamma_1} \frac{1}{w-1} e^{Nf(t_0(w),w)} e^{N(b\ln\tau - \tau)} \frac{\mathrm{d}t}{\mathrm{d}\tau} \mathrm{d}w \mathrm{d}\tau. \end{aligned} \tag{3.6}$$

Next, let us examine the mapping (2.12). We note that this mapping still works in the present case; the only difference is that for fixed $n/N \in (0,1)$, the constant η in the mapping (2.12) is bounded away from 0, whereas when $n/N \to 0$, η approaches 0. More precisely, we have $\eta \sim -b^2$. This can roughly be seen from the saddle points (2.9) and (2.11), together with their correspondence relation (2.13) under the mapping $w \to u$ given in (2.12). To prove this rigorously, we give the following lemma.

Lemma 1. *Let η be the constant defined in the mapping (2.12). If $a/b^2 = o(1)$ (or, equivalently, $xN/n^2 = o(1)$), then we have*

$$\eta = -b^2 + o(b^2) \qquad \text{as } b \to 0. \tag{3.7}$$

Proof. From (2.5), we have for any fixed w

$$t_0(w) = \frac{1}{1+b} - \frac{b^2(1-w)}{1+b} - \frac{b^4 w(w-1)^2}{1+b} + O\left(b^6 w^2 (w-1)^3\right) \qquad \text{as } b \to 0, \tag{3.8}$$

where the term $O\left(b^4 w(w-1)^2\right)$ holds uniformly when either w or $1-w$ is small. If $a/b^2 \to 0$, from (2.9) we have $w_- = 1 - w_+ \to 0^+$ and both $w_\pm$ bounded. Thus,

$$\begin{aligned} t_0(w_-) &= 1 - b + b^2(1-b)w_- + O\left(b^4 w_-^2\right), \\ t_0(w_+) &= \frac{1}{1+b} - \frac{b^2}{1+b} w_- + O\left(b^4 w_-^2\right). \end{aligned} \tag{3.9}$$

To obtain η, we use (2.12) and (2.13). First, substituting (3.9) in (2.2) gives

$$f(t_0(w_+), w_+) - f(t_0(w_-), w_-) = 2a \ln \frac{1-w_-}{w_-} - b^2 + O\left(b^3\right). \tag{3.10}$$

Here, we have made use of the fact that if $w_\pm$ lie on the upper edge of the cut along the interval $0 < w < 1$, then we have $\ln(w_\pm - 1) = \ln(1 - w_\pm) + \pi i$. Similarly, if $w_\pm$ lie on the lower edge of the cut, then $\ln(w_\pm - 1) = \ln(1 - w_\pm) - \pi i$. Next, we have from (2.10) and (2.11)

$$\psi(u_-) - \psi(u_+) = 2a \ln \frac{1-u_+}{u_+} + \eta \sqrt{1 + \frac{4a}{\eta}}. \tag{3.11}$$

Note that $u_+ + u_- = 1$ and $\eta < 0$. Formula (3.7) now follows from a combination of (2.12), (2.13), (3.10) and (3.11). □

What Lemma 1 says is that if $a/b^2 \to 0$ and $x = O(1)$ or $x \gg O(1)$, then we have $|\eta| N \sim b^2 N = x b^2/a \to \infty$, i.e., ηN is large.

Coupling (3.6) and (2.12), we have

$$\begin{aligned} t_n(x, N+1) =& \frac{(-1)^{n+1}\Gamma(n+N+2) e^n b^{-n} e^{N\gamma}}{2\pi i \Gamma(n+1)\Gamma(N-n+1)} \\ & \times \int_0^{+\infty} \int_{\gamma_1} \frac{h(u,\tau)}{u-1} e^{N\psi(u)} e^{N(b\ln\tau - \tau)} \mathrm{d}w \mathrm{d}\tau, \end{aligned} \tag{3.12}$$

where

$$h(u,\tau) = \frac{u-1}{w-1} \frac{\mathrm{d}w}{\mathrm{d}u} \frac{\mathrm{d}t}{\mathrm{d}\tau}, \tag{3.13}$$

$\mathrm{d}w/\mathrm{d}u$ is given by (2.14) and (2.15) and $\mathrm{d}t/\mathrm{d}\tau$ is given by (3.5). Following the same integration-by-parts procedure outlined in Section 2, we let $h_0(u,\tau) = h(u,\tau)$ in (3.13), and define $h_l(u,\tau)$ recursively by (2.18), (2.19) and (2.20). The final result is

$$t_n(x,N+1) = \frac{(-1)^{n+1}\Gamma(n+N+2)e^n b^{-n}e^{N\gamma}}{\Gamma(N-n+1)N^{n+1}}$$

$$\times\left[\mathbf{M}(aN+1,1,\eta N)\sum_{l=0}^{p-1}\frac{c_l}{(\eta N)^l}+\mathbf{M}'(aN+1,1,\eta N)\sum_{l=0}^{p-1}\frac{d_l}{(\eta N)^l}+\varepsilon_p\right], \tag{3.14}$$

where

$$c_l = \frac{N^{n+1}}{\Gamma(n+1)}\int_0^\infty a_l(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^\infty a_{l,m}\frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m}, \tag{3.15}$$

$$d_l = \frac{N^{n+1}}{\Gamma(n+1)}\int_0^\infty b_l(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^\infty b_{l,m}\frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m} \tag{3.16}$$

and

$$\varepsilon_p = \frac{N^{n+1}}{2\pi i\Gamma(n+1)(\eta N)^p}\int_0^{+\infty}\int_{\gamma_1}\frac{h_p(u,\tau)}{u-1}e^{N\psi(u)}e^{N(b\ln\tau-\tau)}\mathrm{d}w\mathrm{d}\tau. \tag{3.17}$$

Note that since $b = n/N \to 0$, the series in (3.15) and (3.16) are asymptotic. Moreover, since ηN is large in this case, (3.14) is indeed a compound asymptotic expansion as $N \to \infty$.

Now, we consider the subcase in which $a/b^2 \to 0$ and $x = o(1)$. In this case, the expansion in (3.14) is still asymptotic as long as $|\eta|N \sim b^2 N \to \infty$ or, equivalently, $n \gg O(\sqrt{N})$. If $n = O(\sqrt{N})$, then the series in (1.2) is itself an asymptotic expansion as $N \to \infty$. This completes our discussion of all three cases listed under the condition "xN/n^2 small" in Table 1.

4 AIRY-TYPE EXPANSION

For the case $a/b^2 \in [\gamma, M]$, where γ is a small positive number and M is a large positive number, the saddle points $w_\pm$ in (2.9) are bounded away from the singularities $w = 0$, $w = 1$ and $w = \infty$, and coalesce with each other when $a/b^2 \to \frac{1}{4(1-a)}$. Therefore, to derive an asymptotic expansion uniformly for $a/b^2 \in [\gamma, M]$, we need the cubic transformation (see (4.6) below) introduced by Chester, Friedman and Ursell [2]. The resulting expansion is in terms of the Airy function $\mathrm{Ai}(\cdot)$.

Following the same argument as in Section 3, we again use the mapping from $t \to \tau$ in (3.2), and start with the integral representation (3.6)

$$t_n(x,N+1) = \frac{(-1)^{n+1}}{2\pi i}\frac{\Gamma(n+N+2)e^n b^{-n}}{\Gamma(n+1)\Gamma(N-n+1)} \times\int_0^{+\infty}\int_{\gamma_1}\frac{1}{w-1}e^{Nf(t_0(w),w)}e^{N(b\ln\tau-\tau)}\frac{\mathrm{d}t}{\mathrm{d}\tau}\mathrm{d}w\mathrm{d}\tau, \tag{4.1}$$

where the integration path is described in the line below (2.2). To proceed further, we divide γ_1 into two parts, and denote the part in the upper half of the plane by $\gamma^{(+)}$, and

the other part in the lower half of the plane by $\gamma^{(-)}$. Recall from (2.2) that the phase function in (4.1) is given by

$$\begin{aligned} f(t_0(w),w) =& b\ln(1-t_0(w)) + (1-b)\ln t_0(w) + a\ln w \\ & - a\ln(w-1) + b\ln[1-(1-t_0(w))w], \end{aligned} \tag{4.2}$$

which has a cut $(-\infty, 1]$ in w-plane. Put

$$\begin{aligned} \overline{f}(t_0(w),w) =& b\ln(1-t_0(w)) + (1-b)\ln t_0(w) + a\ln w \\ & - a\ln(1-w) + b\ln[1-(1-t_0(w))w], \end{aligned} \tag{4.3}$$

which has cuts $(-\infty, 0]$ and $[1, \infty)$. From (4.2) and (4.3), we have

$$f(t_0(w),w) = \overline{f}(t_0(w),w) + a\pi i \tag{4.4}$$

for $w \in \gamma^{(+)}$, and

$$f(t_0(w),w) = \overline{f}(t_0(w),w) - a\pi i \tag{4.5}$$

for $w \in \gamma^{(-)}$. We now make the standard transformation

$$\overline{f}(t_0(w),w) = \frac{1}{3}u^3 - \zeta u + A, \tag{4.6}$$

with the correspondence between the critical points of the two sides prescribed by

$$w_+ \leftrightarrow \sqrt{\zeta}, \qquad w_- \leftrightarrow -\sqrt{\zeta}. \tag{4.7}$$

If $\zeta > 0$, then $\pm\sqrt{\zeta}$ are both real; if $\zeta < 0$, then $\pm\sqrt{\zeta}$ are complex conjugates and purely imaginary. The values of A and ζ can be obtained by using (4.6) and (4.7). We also have

$$\begin{aligned} &\frac{\mathrm{d}w}{\mathrm{d}u} = \frac{(u-\sqrt{\zeta})(u+\sqrt{\zeta})w(1-w)\left[(2a-1) - \sqrt{1+4b^2w^2-4b^2w}\right]}{-2b^2(w-w_+)(w-w_-)}, \quad u \neq \pm\sqrt{\zeta}, \\ &\frac{\mathrm{d}w}{\mathrm{d}u} = \left\{\frac{2\sqrt{\zeta}a(1-2a)(1-a)}{b^3\sqrt{b^2-4a+4a^2}}\right\}^{1/2}, \qquad u = \pm\sqrt{\zeta}, \end{aligned} \tag{4.8}$$

where we have again used L'Hôpital's rule for $u = \pm\sqrt{\zeta}$. Let us first consider the case $\zeta < 0$, and deform the image of the contour $\gamma^{(+)}$ under the mapping $w \to u$ defined in (4.6) to the steepest descent path of $\frac{1}{3}u^3 - \zeta u + A$ in the u-plane which passes through $\sqrt{\zeta}$. We denote the path by C_2. Similarly, we deform the image of the contour $\gamma^{(-)}$, and denote the steepest descent path passing through $-\sqrt{\zeta}$ by C_3; see Figure 1. Next, we consider the case $\zeta > 0$. Note that here the saddle points $\pm\sqrt{\zeta}$ are real, and that the contours C_2 and C_3 pass through both of them; see Figure 2. Clearly, in both cases, C_3 is the reflection of C_2 with respect to the real axis in the u-plane.

Let C_1 denote the dotted curve shown in Figures 1 and 2, and recall the identities $e^{iaN\pi} = \cos aN\pi + i\sin aN\pi$ and $e^{-iaN\pi} = \cos aN\pi - i\sin aN\pi$. A combination of (4.1), (4.4), (4.5) and (4.6) then gives

$$\begin{aligned} t_n(x,N+1) =& \frac{(-1)^n\Gamma(n+N+2)e^n b^{-n}e^{NA}}{\Gamma(n+1)\Gamma(N-n+1)} \\ & \times \left\{ \frac{\cos aN\pi}{2\pi i}\int_0^{+\infty}\int_{C_1} h_0(u,\tau)e^{N\left(\frac{1}{3}u^3-\zeta u\right)}e^{N(b\ln\tau-\tau)}\mathrm{d}u\mathrm{d}\tau \right. \\ & \left. + \frac{\sin aN\pi}{2\pi}\int_0^{+\infty}\int_{C_3-C_2} h_0(u,\tau)e^{N\left(\frac{1}{3}u^3-\zeta u\right)}e^{N(b\ln\tau-\tau)}\mathrm{d}u\mathrm{d}\tau \right\}, \end{aligned} \tag{4.9}$$

where

$$h_0(u,\tau)=\frac{1}{w-1}\frac{\mathrm{d}t}{\mathrm{d}\tau}\frac{\mathrm{d}w}{\mathrm{d}u}, \tag{4.10}$$

$\mathrm{d}t/\mathrm{d}\tau$, $\mathrm{d}w/\mathrm{d}u$ are given respectively by (3.5) and (4.8).




Figure 1: C_2 and C_3 ($\zeta<0$).




Figure 2: C_2 and C_3 ($\zeta>0$).

4.1 When ζ is bounded

Following the standard integration-by-parts procedure [11, p.368], we define recursively

$$h_l(u,\tau) = a_l(\tau)+b_l(\tau)u-(u^2-\zeta)g_l(u,\tau), \tag{4.11}$$

$$h_{l+1}(u,\tau) = \frac{\partial g_l(u,\tau)}{\partial u}. \tag{4.12}$$

Furthermore, expand

$$a_l(\tau)=\sum_{j=0}^{\infty}a_{l,j}\tau^j, \qquad b_l(\tau)=\sum_{j=0}^{\infty}b_{l,j}\tau^j. \tag{4.13}$$

From (4.9), (4.11), (4.12) and (4.13), it follows

$$\begin{aligned} t_n(x,N+1)=&\frac{(-1)^n\Gamma(n+N+2)e^n b^{-n}e^{NA}}{\Gamma(N-n+1)N^{n+1}}\\ &\times\left[\cos aN\pi\left(\mathrm{Ai}(N^{2/3}\zeta)\sum_{l=0}^{p-1}\frac{c_l}{N^{l+1/3}}-\mathrm{Ai}'(N^{2/3}\zeta)\sum_{l=0}^{p-1}\frac{d_l}{N^{l+2/3}}\right)\right.\\ &\left.+\sin aN\pi\left(\mathrm{Bi}(N^{2/3}\zeta)\sum_{l=0}^{p-1}\frac{c_l}{N^{l+1/3}}-\mathrm{Bi}'(N^{2/3}\zeta)\sum_{l=0}^{p-1}\frac{d_l}{N^{l+2/3}}\right)+\varepsilon_p\right], \end{aligned} \tag{4.14}$$

where

$$\begin{aligned} \varepsilon_p=&\frac{\cos(aN\pi)N^{n+1}}{2\pi iN^p\Gamma(n+1)}\int_0^{+\infty}\int_{C_1}h_p(u,\tau)e^{N\left(\frac{1}{3}u^3-\zeta u\right)}e^{N(b\ln\tau-\tau)}\mathrm{d}u\mathrm{d}\tau\\ &+\frac{\sin(aN\pi)N^{n+1}}{2\pi iN^p\Gamma(n+1)}\int_0^{+\infty}\int_{C_3-C_2}h_p(u,\tau)e^{N\left(\frac{1}{3}u^3-\zeta u\right)}e^{N(b\ln\tau-\tau)}\mathrm{d}u\mathrm{d}\tau, \end{aligned} \tag{4.15}$$

and the coefficients are given by

$$
\begin{aligned}
c_l &= \frac{N^{n+1}}{\Gamma(n+1)} \int_0^\infty a_l(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^{\infty} a_{l,m}\frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m},\\
d_l &= \frac{N^{n+1}}{\Gamma(n+1)} \int_0^\infty b_l(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^{\infty} b_{l,m}\frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m}.
\end{aligned} \tag{4.16}
$$

In the present case, $a/b^2 \in [\delta, M]$ or, equivalently, $w_\pm$ are bounded. Thus ζ is also bounded in view of (4.7). Therefore, it is easy to prove that (4.14) is asymptotic; for details, see [11, p.371-372].

4.2 When $\zeta \to -\infty$

When $a/b^2 \to \infty$, by (2.9) the saddle points $w_\pm$ approach $\frac{1}{2} \pm i\infty$, respectively, along the line $\mathrm{Re}\ w = \frac{1}{2}$. In view of (4.7), this is equivalent to saying that $\zeta \to -\infty$. To prove that (4.14) is also asymptotic in this case, we rewrite the expansion in (4.14) as

$$
\begin{aligned}
t_n(x, N+1) =& \frac{(-1)^n\Gamma(n+N+2)e^n b^{-n}e^{NA}}{\Gamma(N-n+1)N^{n+1}}\\
&\times \left[\cos aN\pi \left(\frac{\mathrm{Ai}(N^{2/3}\zeta)}{N^{1/3}}\sum_{l=0}^{p-1}\frac{\widetilde{c}_l}{(aN)^l} - \frac{\mathrm{Ai}'(N^{2/3}\zeta)}{N^{2/3}}\sum_{l=0}^{p-1}\frac{\widetilde{d}_l}{(aN)^l}\right)\right.\\
&\left. + \sin aN\pi \left(\frac{\mathrm{Bi}(N^{2/3}\zeta)}{N^{1/3}}\sum_{l=0}^{p-1}\frac{\widetilde{c}_l}{(aN)^l} - \frac{\mathrm{Bi}'(N^{2/3}\zeta)}{N^{2/3}}\sum_{l=0}^{p-1}\frac{\widetilde{d}_l}{(aN)^l}\right) + \varepsilon_p\right],
\end{aligned} \tag{4.17}
$$

where ε_p is the same as in (4.15) and

$$
\widetilde{c}_l = a^l c_l \quad \text{and} \quad \widetilde{d}_l = a^l d_l. \tag{4.18}
$$

Thus, it is sufficient to first prove the boundedness of the coefficients $\widetilde{c}_l$ and $\widetilde{d}_l$, and then establish the asymptotic nature of the error term in (4.17), namely, to prove that there exist positive constants A_p, A'_p, B_p and B'_p such that

$$
\begin{aligned}
\left|\varepsilon_p^1\right| \le& \frac{A_p}{(aN)^p}\frac{\widetilde{\mathrm{Ai}}(N^{2/3}\zeta)}{N^{1/3}} + \frac{A'_p}{(aN)^p}\frac{\widetilde{\mathrm{Ai}}'(N^{2/3}\zeta)}{N^{2/3}},\\
\left|\varepsilon_p^2\right| \le& \frac{B_p}{(aN)^p}\frac{\widetilde{\mathrm{Bi}}(N^{2/3}\zeta)}{N^{1/3}} + \frac{B'_p}{(aN)^p}\frac{\widetilde{\mathrm{Bi}}'(N^{2/3}\zeta)}{N^{2/3}},
\end{aligned} \tag{4.19}
$$

where

$$
\begin{aligned}
\varepsilon_p^1 =& \frac{\cos(aN\pi)N^{n+1}}{2\pi i N^p\Gamma(n+1)}\int_0^{+\infty}\int_{C_1} h_p(u,\tau)e^{N\left(\frac{1}{3}u^3-\zeta u\right)}e^{N(b\ln\tau-\tau)}\mathrm{d}u\mathrm{d}\tau,\\
\varepsilon_p^2 =& \frac{\sin(aN\pi)N^{n+1}}{2\pi i N^p\Gamma(n+1)}\int_0^{+\infty}\int_{C_3-C_2} h_p(u,\tau)e^{N\left(\frac{1}{3}u^3-\zeta u\right)}e^{N(b\ln\tau-\tau)}\mathrm{d}u\mathrm{d}\tau,
\end{aligned} \tag{4.20}
$$

and $\widetilde{\mathrm{Ai}}(z) = \widetilde{\mathrm{Bi}}(z) = \left[\mathrm{Ai}^2(z) + \mathrm{Bi}^2(z)\right]^{1/2}$ for $z < 0$. Note that in the present case, $\zeta < 0$.

From (4.8), we recall that $\mathrm{d}w/\mathrm{d}u$ depends on ζ. The value of ζ is obtained by solving the two equations gotten from (4.6) with w and u replaced respectively by $w_\pm$ and $\pm\sqrt{\zeta}$; see [11, p.367]. Thus, ζ and hence $\mathrm{d}w/\mathrm{d}u$ both depend on the parameters a and b in (1.3). As a and b approach zero, ζ may tend to infinity. When the saddle points $u_\pm = \pm\sqrt{\zeta}$ are bounded, the mapping $w \to u$ defined by the cubic transformation (4.6) is analytic, and its derivative $\mathrm{d}w/\mathrm{d}u$ is bounded. However, in the present case, the saddle points $u_\pm = \pm\sqrt{-\zeta}i$ go to infinity as $\zeta \to -\infty$. Hence, the coefficients $a_l(\tau)$, $b_l(\tau)$ and the function $h_l(u,\tau)$ given recursively in (4.11) and (4.12) may all blow up as $\zeta \to -\infty$, where $h_0(u,\tau)$ is defined by (4.10). Since the coefficients c_l and d_l in (4.14) are related to $a_l(\tau)$ and $b_l(\tau)$ via (4.16) and (4.13), to prove that the expansion in (4.17) is asymptotic when $\zeta \to -\infty$, we must first give estimates for the coefficient functions $a_l(\tau)$ and $b_l(\tau)$. To this end, we shall adopt a method introduced by Olde Daalhuis and Temme [5]. To begin with, we define

$$A_0(u,\zeta) = \frac{u}{u^2-\zeta}, \qquad B_0(u,\zeta) = \frac{1}{u^2-\zeta}. \tag{4.21}$$

Using (4.11) and the Cauchy residue theorem, it can be verified that

$$\begin{aligned} a_0(\tau) =& \frac{1}{2\pi i}\int_\Gamma h_0(u,\tau)A_0(u,\zeta)\mathrm{d}u, \\ b_0(\tau) =& \frac{1}{2\pi i}\int_\Gamma h_0(u,\tau)B_0(u,\zeta)\mathrm{d}u, \end{aligned} \tag{4.22}$$

where Γ is the contour consisting of two circles, centering at $\pm\sqrt{-\zeta}i$, both with radius R, where R could be as large as possible until the circles reach the singularities of $h_0(u,\tau)$ in the u-plane.

We note that $w = w_\pm$ are removable singularities of $\mathrm{d}w/\mathrm{d}u$. However, $\mathrm{d}w/\mathrm{d}u$ blows up at the points $w = w_\pm e^{2k\pi i}$, where $k \neq 0$ is any integer; see (4.8). To find their image points in the u-plane under the mapping (4.6), we take $w_k = w_+ e^{2k\pi i}$ as an example and denote its image point by u_k. The image point of $w_- e^{2k\pi i}$ can be treated in a similar manner. By (4.6), we have

$$\overline{f}(t_0(w_k), w_k) = \frac{1}{3}u_k^3 - \zeta u_k + A. \tag{4.23}$$

From (4.6) and (4.7), we also have

$$\overline{f}(t_0(w_+), w_+) = \frac{1}{3}u_+^3 - \zeta u_+ + A. \tag{4.24}$$

Subtracting (4.24) from (4.23) gives

$$2k\pi a i = i(-\zeta)^{3/2}\left(-\frac{1}{3}z_k^3 - z_k^2\right). \tag{4.25}$$

Since $0 < a \leq \frac{1}{2}$ and $-\zeta \gg O(1)$, from (4.25) it follows that $|z_k| \sim \sqrt{2k\pi}a^{1/2}(-\zeta)^{-3/4}$. Therefore, there exists a constant $0 < c_0 < 1$, independent of a, b and ζ, such that the interior of the two circles with centres at $u = \pm\sqrt{-\zeta}i$ and radius $R = c_0\sqrt{a}(-\zeta)^{-1/4}$ is

free of the singularities of $h_0(u,\tau)$ in the u-plane. Since $h_0(u,\tau)$ is now analytic inside the contour Γ, there exists a constant c_h such that

$$|h_0(u,\tau)| \le c_h h(\tau) \tag{4.26}$$

for u in the domain enclosed by the contour Γ, where $h(\tau)$ denotes the maximum of the two functions $\left|h_0(\pm\sqrt{-\zeta}i,\tau)\right|$. Note that as functions of τ, $h_0(\pm\sqrt{-\zeta}i,\tau)$ are analytic in the neighbourhood of steepest descent path in the τ-plane.

We further introduce rational functions A_k and B_k, $k = 0, 1, 2...$, defined recursively by

$$\begin{aligned} A_{k+1}(u,\zeta) =& \frac{1}{u^2-\zeta}\frac{\mathrm{d}}{\mathrm{d}u}A_k(u,\zeta), \\ B_{k+1}(u,\zeta) =& \frac{1}{u^2-\zeta}\frac{\mathrm{d}}{\mathrm{d}u}B_k(u,\zeta), \end{aligned} \tag{4.27}$$

where A_0 and B_0 are given in (4.21). By induction, we can show that A_k and B_k are expressible as

$$\begin{aligned} A_k(u,\zeta) =& \sum_{i=0}^{[(k+1)/2]} \frac{c^1_{k,i}u^{k+1-2i}}{\left(u^2-\zeta\right)^{2k+1-i}}, \\ B_k(u,\zeta) =& \sum_{i=0}^{[k/2]} \frac{c^2_{k,i}u^{k-2i}}{\left(u^2-\zeta\right)^{2k-i}}, \end{aligned} \tag{4.28}$$

where $c^1_{k,i}$ and $c^2_{k,i}$ are constants independent of u and ζ. As in (4.22), by Cauchy's theorem, we have from equations (4.11) and (4.12)

$$\begin{aligned} a_k(\tau) =& \frac{1}{2\pi i}\int_\Gamma h_k(u,\tau)A_0(u,\zeta)\mathrm{d}u \\ =& \frac{1}{2\pi i}\int_\Gamma h_{k-1}(u,\tau)A_1(u,\zeta)\mathrm{d}u - \frac{1}{2\pi i}\int_\Gamma \left(a_{k-1}(\tau)+b_{k-1}(\tau)u\right)A_1(u,\zeta)\mathrm{d}u \\ =& \frac{1}{2\pi i}\int_\Gamma h_{k-1}(u,\tau)A_1(u,\zeta)\mathrm{d}u \\ &\vdots \\ =& \frac{1}{2\pi i}\int_\Gamma h_0(u,\tau)A_k(u,\zeta)\mathrm{d}u, \end{aligned} \tag{4.29}$$

where we have used integration-by-parts to derive the second equality. The second term in the second equality vanishes because $\left(a_{k-1}(\tau)+b_{k-1}(\tau)u\right)A_1(u,\zeta)$ is $O\left(u^{-2}\right)$ as $|u|\to\infty$ and all poles of that function lie inside Γ; see (4.28). Similarly, we also have

$$b_k(\tau) = \frac{1}{2\pi i}\int_\Gamma h_0(u,\tau)B_k(u,\zeta)\mathrm{d}u. \tag{4.30}$$

Using (4.28), it is easy to obtain the estimates

$$|A_k(u,\zeta)| \le C_k a^{-k-1/2}(-\zeta)^{1/4} \quad \text{and} \quad |B_k(u,\zeta)| \le C_k a^{-k}, \tag{4.31}$$

for u on and inside the contour Γ and $-\zeta \to \infty$. Here and thereafter, C_k is used as a generic symbol for constants independent of u, ζ, a and b. Substituting (4.31) into (4.29) and (4.30) gives

$$|a_k(\tau)| \le C_k h(\tau) a^{-k}, \tag{4.32}$$

and

$$|b_k(\tau)| \le C_k h(\tau) a^{-k+1/2} (-\zeta)^{-1/4}. \tag{4.33}$$

Therefore, $a^l a_l(\tau)$ and $a^l b_l(\tau)$ are both bounded for τ in the neighbourhood of steepest descent path in the τ-plane, and of course for τ in the neighbourhood of $\tau = 0$. Thus we have the boundedness of the coefficients $\widetilde{c}_l$ and $\widetilde{d}_l$.

To estimate $h_k(u, \tau)$, we use the rational functions $R_k(u, w, \zeta)$, $k = 0, 1, 2...$, defined recursively by

$$\begin{aligned} R_0(u, w, \zeta) =& \frac{1}{u - w}, \\ R_{k+1}(u, w, \zeta) =& \frac{1}{u^2 - \zeta} \frac{\mathrm{d}}{\mathrm{d}u} R_k(u, w, \zeta). \end{aligned} \tag{4.34}$$

These functions were also introduced by Olde Daalhuis and Temme [5]. They showed by induction that $R_k(u, w, \zeta)$ can be written as

$$R_k(u, w, \zeta) = \sum_{i=0}^{k-1} \sum_{j=0}^{\min\{i, k-1-i\}} \frac{C_{ij} u^{i-j}}{(u - w)^{k+1-i-j} \left(u^2 - \zeta\right)^{k+i}}, \qquad k = 1, 2..., \tag{4.35}$$

where C_{ij} do not depend on u, w and ζ. Similar to (4.29), we have

$$h_k(w, \tau) = \frac{1}{2\pi i} \int_\Gamma h_0(u, \tau) R_k(u, w, \zeta) \mathrm{d}u, \tag{4.36}$$

where Γ is the same contour used in (4.29) and w lies inside two disks centered at $\pm\sqrt{-\zeta} i$ and with radius $\frac{1}{2} c_0 \sqrt{a} (-\zeta)^{-1/4}$. It is easy to verify from (4.35) that

$$|R_k(u, w, \zeta)| \le C_k a^{-k-1/2} (-\zeta)^{1/4}, \tag{4.37}$$

and from (4.36) that

$$|h_k(w, \tau)| \le C_k a^{-k} h(\tau). \tag{4.38}$$

Substituting (4.38) into (4.20) gives (4.19); for details, see [5, p.311-312]. Note that to make the expansion (4.17) asymptotic, we require $x = aN$ to be large.

5 BESSEL-TYPE EXPANSION

In the case of Hahn Polynomials $Q_n(x; \alpha, \beta, N)$ given in (1.1), Sharapodinov [10] has given an asymptotic formula involving Jacobi polynomials when the parameters satisfy α, $\beta \ge \frac{1}{2}$ and $2 \le n \le c\sqrt{N}$, where c is a positive constant. The values of the variable

x are also required to be large; more precisely, $x \geq \varepsilon N$ and $\varepsilon > 0$ is a small number. Although discrete Chebyshev polynomials $t_n(x, N)$ given in (1.2) is a special case of the Hahn polynomials, the values of the parameters are $\alpha = \beta = 0$; that is, Sharapodinov's result does not include our case. However, since the leading term in the uniform asymptotic expansion of the Jacobi polynomials is a Bessel function (see [7, p.451]), the work of Sharapodinov did inspire us to look for an asymptotic expansion for $t_n(x, N+1)$ involving Bessel functions, when the parameters a and b in (1.3) satisfy $a/b^2 \to \infty$ and the variable x is large. Our method differs completely from that of Sharapodinov.

Returning to (2.1), and making the change of variable $v = 1/w$, we have

$$t_n(x, N+1) = \frac{(-1)^n}{2\pi i} \frac{\Gamma(n+N+2)}{\Gamma(n+1)\Gamma(N-n+1)} \int_0^1 \int_{\gamma_3} \frac{1}{v(1-v)} e^{N\widehat{f}(t,v)} \mathrm{d}v\mathrm{d}t, \tag{5.1}$$

where

$$\widehat{f}(t, v) = b \ln(1-t) + (1-b) \ln t - a \ln(1-v) - b \ln v + b \ln[t+v-1]; \tag{5.2}$$

the curve γ_3 starts at $v = +\infty$, runs along the lower edge of the positive real line towards $v = 1$, encircles the point $v = 1$ in the clockwise direction and returns to $+\infty$ along the upper edge of the positive real line. Note that since $e^{Nb \ln v} = v^n$, there is no need to have a cut from the origin to infinity.

The saddle point of $\widehat{f}(t, v)$ in the t-plane is given by

$$t_0(v) = \frac{2 - v + \sqrt{4b^2 - 4b^2 v + v^2}}{2(1+b)}; \tag{5.3}$$

cf.(2.5), where the v-plane is cut along two line segments joining 0 to the two conjugate points $2b^2 \pm 2b\sqrt{1-b^2}i$, and the branch of the square root is chosen so that $\sqrt{4b^2 - 4b^2 v + v^2} \sim v$ as $v \to \infty$. From (5.3), we have

$$\begin{aligned} t_0(v) &= 1 + O(b) \qquad \text{as } |v| \sim cb, \\ t_0(v) &\sim 1 - b \qquad\quad \text{as } v \gg b, \end{aligned} \tag{5.4}$$

where c is a positive constant, and for $v \ll b$,

$$\begin{aligned} t_0(v) &\sim 1 - \frac{1}{2} v \qquad \text{as } v \to 0^+, \\ t_0(v) &\to 1 - 2b \qquad \text{as } v \to 0^-, \end{aligned} \tag{5.5}$$

where 0^+ and 0^- mean limits approaching 0 from the right-hand side of the cut and the left-hand side of the cut, respectively. Moreover, easy calculation shows that for any v, $t_0(v)$ is not a real number on the cut $(1, \infty)$ in the t-plane. Following the same argument given prior to (3.2), we introduce the mapping

$$\widehat{f}(t, v) = b \ln \tau - \tau + A \tag{5.6}$$

with the correspondence between the saddle points $t = t_0(v)$ and $\tau = b$ given by

$$\tau(t_0(v)) = b. \tag{5.7}$$

Coupling (5.6) and (5.7) yields

$$A = \widehat{f}(t_0(v), v) - b\ln b + b. \tag{5.8}$$

The zeros of $\partial \widehat{f}(t,v)/\partial t = 0$ are given by

$$t_0^{\pm}(v) = \frac{2 - v \pm \sqrt{4b^2 - 4b^2 v + v^2}}{2(1+b)}, \tag{5.9}$$

where $t_0^+(v)$ is the relavant saddle point $t_0(v)$ given in (5.3). By straightforward calculation, we have

$$\begin{aligned} \frac{\mathrm{d}t}{\mathrm{d}\tau} &= \frac{b/\tau - 1}{\partial \widehat{f}(t,v)/\partial t} = \frac{(\tau - b)t(1-t)(t+v-1)}{\tau(1+b)(t - t_0^+(v))(t - t_0^-(v))}, \qquad \tau \neq b, \\ \frac{\mathrm{d}t}{\mathrm{d}\tau} &= \left\{ \frac{t_0^+(v)(1 - t_0^+(v))(t_0^+(v) + v - 1)}{b\sqrt{4b^2 - 4b^2 v + v^2}} \right\}^{1/2}, \qquad \tau = b. \end{aligned} \tag{5.10}$$

Note that we have $\tau = 0$ when $t = 1$, and $\tau = +\infty$ when $t = 0$. For any fixed v, we can deform the original interval of integration $0 \le t \le 1$ into a steepest descent path Γ_v, passing through $t_0(v)$. Also note that $t_0^-(v)$ is not on Γ_v, unless $0 < v < 1$. Moreover, if $0 < v < 1$, then $0 < t_0^-(v) < t_0^+(v) < 1$ and Γ_v is the real interval $[0, 1]$. In our case, there is only one point on the path γ_3 in the v-plane (see (5.1)), where it crosses the real line. Let us denote this point by v_0. When $v = v_0$, the mapping which we have introduced in (5.6) becomes singular at the point $t_0^-(v)$, since $\mathrm{d}t/\mathrm{d}\tau$ in (5.10) blows up. However for this particular case, we only need to slightly modify the path by replacing part of the original path near this point by a small half circle as shown in Figure 3. The




Figure 3: The indentation of the integration path in the t-plane for $0 < v < 1$.

corresponding integration path in the τ-plane following the mapping (5.6) also needs to be modified. However, this small modification on the integration path will not affect the following argument and calculation. With this in mind, we will simply ignore this particular case, and proceed with the assumption that the integration path in the τ-plane is always $[0, +\infty)$ for all w, and that $\mathrm{d}t/\mathrm{d}\tau$ will not blow up in the neighbourhood of the path.

Thus, from (5.1) and (5.6), we have

$$\begin{aligned} t_n(x, N+1) =& \frac{(-1)^{n+1}}{2\pi i} \frac{\Gamma(n+N+2)e^n b^{-n}}{\Gamma(n+1)\Gamma(N-n+1)} \\ & \times \int_0^{+\infty} \int_{\gamma_3} \frac{1}{v(1-v)} e^{N\widehat{f}(t_0(v),v)} e^{N(b\ln\tau - \tau)} \frac{\mathrm{d}t}{\mathrm{d}\tau} \mathrm{d}v \mathrm{d}\tau. \end{aligned} \tag{5.11}$$

Here, we rewrite the phase function $\widehat{f}(t_0(v), v)$ as

$$\widehat{f}(t_0(v), v) = (1-b)\ln t_0(v) + b\ln(1-t_0(v)) - a\ln(1-v) - b\ln(\frac{v}{t_0(v)+v-1}). \tag{5.12}$$

Recalling the statements following (5.2) and (5.3), we know that there are only two cuts in the v-plane: one along the infinite interval $[1, +\infty)$ and the other along the bent line joining the conjugate points $2b^2 \pm 2b\sqrt{1-b^2}i$ and passing through the origin. To find the saddle points of $\widehat{f}(t_0(v), v)$, we set

$$\frac{\partial \widehat{f}(t_0(v), v)}{\partial v} = 0,$$

and obtain

$$v_\pm = \frac{b^2 \mp bi\sqrt{4a - 4a^2 - b^2}}{2a(1-a)}. \tag{5.13}$$

Since (5.1) is obtained from (2.1) by making the change of variable $v = 1/w$, (5.13) can also be derived from (2.9). Note that in this case, $b^2/a \to 0$. Furthermore, since $a < \frac{1}{2}$, the quantity inside the square root is positive. Hence, $v_\pm$ are distinct, and approach $v = 0$.

Define

$$g(u) = m(u - \frac{1}{u}),$$

where $m > 0$ is some constant to be determined. The saddle points of $g(u)$ are

$$u_\pm = \pm i. \tag{5.14}$$

Make the transformation

$$\begin{aligned} \widehat{f}(t_0(v), v) &= g(u) + \gamma \\ &= m(u - \frac{1}{u}) + \gamma \end{aligned} \tag{5.15}$$

with

$$u(v_+) = u_-, \qquad u(v_-) = u_+. \tag{5.16}$$

Note the fact that $t_0(v_\pm) = t_\pm$, where $t_\pm$ are given in (2.3) and (2.4). This can be seen from (2.6) and the change of variable $v = 1/w$ that we have made. Thus, substituting (5.16) into (5.15) gives

$$\begin{aligned} \gamma &= \frac{1}{2}\ln\frac{1-b}{1+b} + \frac{1}{2}b\ln\frac{b^2}{1-b^2} \\ &= b\ln b - b + O(b^3) \end{aligned} \tag{5.17}$$

and

$$\begin{aligned} m = -\frac{1}{2}\Bigg\{ &(1-b)\arctan\frac{b\sqrt{4a-4a^2-b^2}}{2-2a-b^2} - a\arctan\frac{b\sqrt{4a-4a^2-b^2}}{2a-2a^2-b^2} \\ &-2b\arctan\frac{\sqrt{4a-4a^2-b^2}}{2+b-2a}\Bigg\} \\ \sim\ & b\arctan\sqrt{\frac{a}{1-a}}, \end{aligned} \tag{5.18}$$

when $b^2/a \to 0$ and $b \to 0$. Moreover, from (5.15) we have

$$\begin{aligned}
&\frac{\mathrm{d}v}{\mathrm{d}u} = \frac{m(u-i)(u+i)v(1-v)\left[(1-2a)v + \sqrt{4b^2 - 4b^2v + v^2}\right]}{2a(1-a)u^2(v-v_+)(v-v_-)}, \qquad u \neq u_\pm, \\
&\left.\frac{\mathrm{d}v}{\mathrm{d}u}\right|_{u=u_\pm} = \left\{\frac{-2m(v_\mp)^2(1-v_\mp)(1-2a)}{b\sqrt{4a-4a^2-b^2}}\right\}^{1/2}, \qquad u = u_\pm.
\end{aligned} \tag{5.19}$$

Here we have made use of the equality $\sqrt{4b^2 - 4b^2v_\mp + v_\mp^2} = (1-2a)v_\mp$.

Coupling (5.11) and (5.15), the integral representation of $t_n(x, N+1)$ becomes

$$\begin{aligned}
t_n(x, N+1) =& \frac{(-1)^{n+1}}{2\pi i}\frac{\Gamma(n+N+2)e^n b^{-n}e^{N\gamma}}{\Gamma(n+1)\Gamma(N-n+1)} \\
&\times \int_0^{+\infty}\int_{\widehat{\gamma}} \frac{h(u,\tau)}{u} e^{N\left[m\left(u-\frac{1}{u}\right)\right]} e^{N(b\ln\tau - \tau)}\mathrm{d}u\mathrm{d}\tau,
\end{aligned} \tag{5.20}$$

where

$$h(u,\tau) = \frac{u}{v(1-v)}\frac{\mathrm{d}t}{\mathrm{d}\tau}\frac{\mathrm{d}v}{\mathrm{d}u}, \tag{5.21}$$

and the contour $\widehat{\gamma}$ starts from $-\infty$, encircles the point $u = 0$ in the counterclockwise direction and returns to $-\infty$. For $l = 0, 1, 2, \cdots$, we define recursively

$$h_l(u,\tau) = a_l(\tau) + \frac{b_l(\tau)}{u} + \left(1 + \frac{1}{u^2}\right) g_l(u,\tau), \tag{5.22}$$

$$h_{l+1}(u,\tau) = -\frac{u}{m}\frac{\mathrm{d}}{\mathrm{d}u}\left\{\frac{g_l(u,\tau)}{u}\right\}, \tag{5.23}$$

where $h_0(u,\tau) = h(u,\tau)$ given in (5.21). Furthermore, we expand $a_l(\tau)$ and $b_l(\tau)$ at $\tau = 0$, and write

$$a_l(\tau) = \sum_{j=0}^{\infty} a_{l,j}\tau^j, \qquad b_l(\tau) = \sum_{j=0}^{\infty} b_{l,j}\tau^j. \tag{5.24}$$

It is easy to see that

$$\begin{aligned}
a_l(\tau) &= \frac{1}{2}\left[h_l(i,\tau) + h_l(-i,\tau)\right], \\
b_l(\tau) &= \frac{i}{2}\left[h_l(i,\tau) - h_l(-i,\tau)\right].
\end{aligned} \tag{5.25}$$

From (5.20), (5.22) and (5.23), we have

$$\begin{aligned}
t_n(x, N+1) =& \frac{(-1)^{n+1}}{2\pi i}\frac{\Gamma(n+N+2)e^n b^{-n}e^{N\gamma}}{\Gamma(n+1)\Gamma(N-n+1)} \\
&\times \int_0^{+\infty}\int_{-\infty}^{(0^+)} \left[a_0(\tau) + \frac{b_0(\tau)}{u} + \left(1 + \frac{1}{u^2}\right) g_0(u,\tau)\right] \frac{e^{Nm\left(u-\frac{1}{u}\right)}}{u}\tau^n e^{-N\tau}\mathrm{d}u\mathrm{d}\tau \\
=& \frac{(-1)^{n+1}\Gamma(n+N+2)e^n b^{-n}e^{N\gamma}}{\Gamma(N-n+1)N^{n+1}}\left[c_0 J_0(2Nm) + d_0 J_1(2Nm) + \varepsilon_1^+\right],
\end{aligned} \tag{5.26}$$

where

$$
\begin{aligned}
c_0 &= \frac{N^{n+1}}{\Gamma(n+1)} \int_0^\infty a_0(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^{\infty} a_{0,m} \frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m}, \\
d_0 &= \frac{N^{n+1}}{\Gamma(n+1)} \int_0^\infty b_0(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^{\infty} b_{0,m} \frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m}
\end{aligned} \tag{5.27}
$$

and

$$
\varepsilon_1^+ = \frac{N^{n+1}}{2\Gamma(n+1)\pi i} \int_0^{+\infty} \int_{-\infty}^{(0^+)} \left(1+\frac{1}{u^2}\right) \frac{g_0(u,\tau)}{u} e^{Nm\left(u-\frac{1}{u}\right)} \tau^n e^{-N\tau}\mathrm{d}u\mathrm{d}\tau. \tag{5.28}
$$

In (5.26), we have made use of the integral representations of Bessel function and Gamma function

$$
J_v(z) = \frac{1}{2\pi i} \int_{-\infty}^{(0^+)} u^{-v-1} \exp\left\{\frac{z}{2}\left(u-\frac{1}{u}\right)\right\} \mathrm{d}u
$$

and

$$
\Gamma(n+1) = N^{n+1} \int_0^{+\infty} e^{-N\tau}\tau^n \mathrm{d}\tau;
$$

see [7, (10.9.19) and (5.9.1)]. Using (5.23) and integration by parts, we can rewrite ε_1^+ in (5.28) as

$$
\varepsilon_1^+ = \frac{N^n}{2\Gamma(n+1)\pi i} \int_0^{+\infty} \int_{-\infty}^{(0^+)} \frac{h_1(u,\tau)}{u} e^{Nm\left(u-\frac{1}{u}\right)} \tau^n e^{-N\tau}\mathrm{d}u\mathrm{d}\tau. \tag{5.29}
$$

Repeating the procedure above, we obtain

$$
\begin{aligned}
t_n(x, N+1) = \frac{(-1)^{n+1}\Gamma(n+N+2)e^n b^{-n} e^{N\gamma}}{\Gamma(N-n+1)N^{n+1}} \Bigg[& J_0(2Nm) \sum_{l=0}^{p-1} \frac{c_l}{N^l} \\
& + J_1(2Nm) \sum_{l=0}^{p-1} \frac{d_l}{N^l} + \varepsilon_p^+ \Bigg],
\end{aligned} \tag{5.30}
$$

where

$$
c_l = \frac{N^{n+1}}{\Gamma(n+1)} \int_0^\infty a_l(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^{\infty} a_{l,m} \frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m}, \tag{5.31}
$$

$$
d_l = \frac{N^{n+1}}{\Gamma(n+1)} \int_0^\infty b_l(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^{\infty} b_{l,m} \frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m}, \tag{5.32}
$$

$$
\varepsilon_p^+ = \frac{N^{n+1-p}}{2\Gamma(n+1)\pi i} \int_0^{+\infty} \int_{-\infty}^{(0^+)} \frac{h_p(u,\tau)}{u} e^{Nm\left(u-\frac{1}{u}\right)} \tau^n e^{-N\tau}\mathrm{d}u\mathrm{d}\tau. \tag{5.33}
$$

Since $h_0(u,\tau)$ involves $\mathrm{d}v/\mathrm{d}u$ and $\mathrm{d}v/\mathrm{d}u$ depends on the parameters a and b in (1.3), the coefficients c_l and d_l in (5.31) and (5.32) also depend on a and b. For an estimate on these coefficients, see (5.57) below.

To facilitate the application of expansion (5.30), we recall that the constants γ and m are explicitly given in (5.17) and (5.18), and note that the leading coefficient can be (asymptotically) calculated by using (5.24) and (5.25). Indeed, we have

$$c_0 \sim a_{0,0} = \frac{1}{2}\left[h_0(i,0) + h(-i,0)\right]$$

and

$$d_0 \sim b_{0,0} = \frac{i}{2}\left[h_0(i,0) - h(-i,0)\right]$$

as $b \to 0$ and $N \to \infty$. Furthermore, equation (5.39) gives

$$h_0(\pm i, 0) = \frac{-(1-a)v_{\mp}}{b}\sqrt{\frac{2(1-2a)m}{b\sqrt{4a-4a^2-b^2}(1-v_{\mp})}}e^{\pm\Delta i}, \tag{5.34}$$

where $\Delta = 2\arctan\sqrt{\frac{a}{1-a}} - \sqrt{\frac{a}{1-a}}$.

5.1 The mapping $v \to u$ in (5.15)

In the case under discussion, $a/b^2 \to \infty$ and $x \to \infty$. Thus, since $0 < a < \frac{1}{2}$, we have $4a - 4a^2 - b^2 > 0$, and the saddle points in (5.13) are complex.

Theorem 1. *When $a \in (0, \frac{1}{2}]$, $b \to 0^+$ and $b^2/a \to 0$, the mapping $v \to u$ defined in (5.15) is one-to-one and analytic for $v \in D_v$ in the v-plane and $u \in D_u$ in the u-plane, where the image of the boundary of D_u in the Z-plane is given in (5.36), and D_v is the image of D_u under the mapping (5.15).*

Proof. As in the cases of Charlier polynomials [9] and Meixner polynomials [3], we introduce an intermediate variable Z defined by

$$\begin{aligned} &b\ln(1-t_0(v)) + (1-b)\ln t_0(v) - a\ln(1-v) - b\ln(\frac{v}{t_0(v)+v-1}) - \gamma \\ &= Z = m\left(u - \frac{1}{u}\right), \end{aligned} \tag{5.35}$$

where $t_0(v)$ is the relevant saddle point given in (5.3).

We consider the upper half of the v-plane. Since the functions $\widehat{f}(t_0(v), v) - \gamma$ and $g(u)$ are both symmetric with respect to the real line, the case for the lower half of the v-plane can be handled in the same manner. To avoid multi-valuedness around the saddle point, we divide the upper half of the v-plane into two parts by using the steepest descent path through v_-, i.e., the point denoted by D; see Figure 4. Call the two parts region I and region III. Note that there are two branch cuts: one along the infinite interval $[1, \infty)$, and the other along the line segment joining $v = 0$ to $v = 2b^2 + i2b\sqrt{1-b^2}$, which was introduced in (5.3). We denoted the point $2b^2 + i2b\sqrt{1-b^2}$ by the letter T in Figure 4.

As v traverses along the boundary $ABCDEFGA$ of region I, the image point Z traverses along the corresponding boundary of a region in the Z-plane; see Figure 5. In Figure 7, we draw the boundary of the region, corresponding to region I, in the u-plane. The image point $Z = \psi(u) + \gamma$ traverses along the boundary of the same region shown




Figure 4: The upper half of the v-plane.




Figure 5: The image of Region I in the Z-plane.

in Figure 5. Note that we have used an arc $GABC$ to avoid the cut OT. The boundary curves EF and $GABC$ in the v-plane are rather arbitrary; for convenience, we choose them to be the ones whose images in the Z-plane are as indicated in Figure 5.

From (5.15), it is readily seen that the mapping $v \to u$ is the composite function of g^{-1} and $\widehat{f}(t_0(v), v) - \gamma$. We have just verified that this mapping is one-to-one on the boundary of region I. By the same argument, one can prove that this mapping is also one-to-one on the boundary of region III. For the image of region III in the intermediate Z-plane, see Figure 6. In the u-plane, we let D_u denote the union of the regions I to IV, with




Figure 6: The image of Region II in the Z-plane.




Figure 7: The upper half of the u-plane.

outer boundary $FEE'KJIVUTN'NM$, and inner boundary $GABCC'HWQ'QRSL$; see Figure 7. Under the mapping $Z = g(u)$, we express the images of parts of the outer and inner boundaries of D_u in the Z-plane as follows:

$$
\begin{array}{rll}
GA: & \text{Re } Z = bM & 0 \le \text{Im } Z \le \theta b\pi; \\
AB: & -bM \le \text{Re } Z \le bM & \text{Im } Z = \theta b\pi; \\
BC \text{ and } C'H: & \text{Re } Z = -bM & 0 \le \text{Im } Z \le \theta b\pi; \\
IJ: & \text{Re } Z = aM & 0 \le \text{Im } Z \le a\pi; \\
JK: & -aM \le \text{Re } Z \le aM & \text{Im } Z = a\pi; \\
KF: & \text{Re } Z = -aM & 0 \le \text{Im } Z \le a\pi,
\end{array}
\tag{5.36}
$$

where $M > 0$ is any fixed constant, and θ is a generic symbol for a constant in $(0,1)$. Furthermore, we denote by D_v the corresponding region of D_u in the v-plane.

By Theorem 1.2.2 of [13, p.12], the mapping $v \to u$ is one-to-one in the interior of both regions I and III in the upper half of the v-plane. As explained earlier, the one-to-one property of this mapping in the lower half of the v-plane can be established by using the symmetry of the functions with respect to the real axis. Note that the only possible singular points of the mapping $v \to u$ in D_v are at $v = v_{\mp}$. Since the images of these points in the u-plane (i.e., $u = \pm i$) are bounded, the mapping is in fact one-to-one and analytic in D_v.

□

5.2 Analyticity of $h_0(u,\tau)$

We now investigate the function $h_0(u,\tau) = h(u,\tau)$ given in (5.21). By (5.10) and (5.19), we have

(5.37)
$$h_0(u,\tau) = \frac{m(\tau-b)t(1-t)(t+v-1)(u-i)(u+i)\left[(1-2a)v+\sqrt{4b^2-4b^2v+v^2}\right]}{2a(1-a)\tau(1+b)(t-t_0^+(v))(t-t_0^-(v))u(v-v_+)(v-v_-)}$$

when $\tau \neq b$ and $u \neq \pm i$, and

(5.38)
$$\begin{aligned} h_0(u,\tau) =& \frac{m(u-i)(u+i)\left[(1-2a)v+\sqrt{4b^2-4b^2v+v^2}\right]}{2a(1-a)u(v-v_+)(v-v_-)} \\ & \times \sqrt{\frac{t_0(v)(1-t_0(v))(t_0(v)+v-1)}{b\sqrt{4b^2-4b^2v+v^2}}} \end{aligned}$$

when $\tau = b$ and $u \neq \pm i$, where $u \in D_u$ and D_u is defined in the previous subsection. Note that $u = 0$ is excluded from D_u. It is easy to see that $(t-1)/\tau$ is bounded by a constant, independent of b and a, as $\tau \to 0$ for $v \in D_v$ and $u \in D_u$. Therefore, $h_0(u,\tau)$ is analytic in τ for τ in the neighbourhood of $[0,\infty)$. Moreover, $h_0(u,\tau)$ is also analytic in u for $u \in D_u$, since the only singularities $u = \pm i$ in D_u are both removable; indeed, we have

(5.39)
$$h_0(u,\tau) = \frac{(\tau-b)t(1-t)(t+v_{\mp}-1)}{\tau(1+b)(t-t_0^+(v_{\mp}))(t-t_0^-(v_{\mp}))}\sqrt{\frac{2(1-2a)m}{(1-v_{\mp})b\sqrt{4a-4a^2-b^2}}}$$

when $\tau \neq b$ and $u = \pm i$, and by the equation following (5.19) we also have

(5.40)
$$h_0(u,\tau) = \sqrt{\frac{2mt_{\mp}(1-t_{\mp})(t_{\mp}+v_{\mp}-1)}{b^2v_{\mp}(1-v_{\mp})\sqrt{4a-4a^2-b^2}}}$$

when $\tau = b$ and $u = \pm i$. To reach (5.40), we have written $u_{\pm}$ as $u_{\pm} = \sqrt{u_{\pm}^2}$ and move $u_{\pm}^2$ inside the square root in $\mathrm{d}v/\mathrm{d}u$ at $u = u_{\pm}$; see (5.19).

For the analysis to be used in the next subsection, we now give an estimate for $h_0(u,0)$ when u lies on the boundaries of D_u, i.e, the outer boundary $FEE'KJIVUTN'NM$ and

the inner boundary $GABCC'HWQ'QRSL$ shown in Figure 7. For convenience, let us denote the outer boundary by C_O and the inner boundary by C_I.

First, we show that for u on the inner boundary C_I,

$$C_1\frac{m}{b} < |u| < C_2\frac{m}{b}, \tag{5.41}$$

where $C_1 < C_2$ and C, C_1, C_2 are used, here and thereafter, as generic symbols for constants independent of u, v, m, a and b. Recall from (5.18) that $m \sim b\arctan\sqrt{\frac{a}{1-a}}$ as $b \to 0$. To prove (5.41), we take the part AB of the inner boundary C_I as an illustration. Using (5.36), we obtain

$$\left| m\left(1 - \frac{1}{|u|^2}\right) \text{Re } u \right| \le bM \quad \text{and} \quad m\left(1 + \frac{1}{|u|^2}\right) \text{Im } u = \theta b\pi. \tag{5.42}$$

Since $|\text{Im } u| \le |u|$, from the equality in (5.42) we have $|u|^2 - \frac{\theta b\pi}{m}|u| + 1 \ge 0$. Therefore,

$$|u| \le \frac{\theta b\pi}{2m} - \sqrt{\left(\frac{\theta b\pi}{2m}\right)^2 - 1} \le \frac{Cm}{b\pi}. \tag{5.43}$$

Rewriting (5.42) gives

$$\frac{m}{|u|^2}\,|\text{Re } u| \le bM + m\,|\text{Re } u| \quad \text{and} \quad \frac{m}{|u|^2}\,|\text{Im } u| \le \theta b\pi + m\,|\text{Im } u|\,, \tag{5.44}$$

which lead to $\frac{m}{|u|} \le bM + \theta b\pi + m|u|$. Substituting (5.43) into the right-hand side of the last inequality, and combining the resulting inequality with (5.43), we obtain (5.41) immediately. Similarly, for u on the outer boundary C_O, we have the estimates

$$C_1\frac{a}{m} < |u| < C_2\frac{a}{m}. \tag{5.45}$$

Hence, it follows from (5.41), (5.45) and (5.37) that

$$|h(u,0)| \le C \qquad \text{for } u \in C_I \cup C_O. \tag{5.46}$$

5.3 Error bounds for the remainder

To prove the asymptotic nature of the expansion in (5.30), we need to give precise estimates for its coefficients c_l, d_l in (5.31) and (5.32), and the error term ε_p^+ given in (5.33), since the derivative $\mathrm{d}v/\mathrm{d}u$ in (5.19) may blow up as $v_\pm$ approach 0, just like the case in Section 4. Because the coefficients c_l and d_l are related to $a_p(\tau)$ and $b_p(\tau)$ by (5.24), (5.31) and (5.32), let us first estimate $a_p(\tau)$ and $b_p(\tau)$. To this end, we define recursively

$$A_0(u) = \frac{u}{1+u^2}, \qquad B_0(u) = -\frac{1}{1+u^2},$$

and

$$\begin{aligned} A_{p+1}(u) &= \frac{1}{m}\left(1 + \frac{1}{u^2}\right)^{-1}\left\{\frac{1}{u} + \frac{\mathrm{d}}{\mathrm{d}u}\right\} A_p(u), \\ B_{p+1}(u) &= \frac{1}{m}\left(1 + \frac{1}{u^2}\right)^{-1}\left\{\frac{1}{u} + \frac{\mathrm{d}}{\mathrm{d}u}\right\} B_p(u) \end{aligned}$$

for $p = 0, 1, 2, ...$ By induction, it can be shown that

$$
\begin{aligned}
A_p(u) &= \left(\frac{1}{2m}\right)^p \left(\frac{1}{u}\right)^{p+1} \left(\frac{u^2}{1+u^2}\right)^{p+1} \sum_{l=0}^{p} c_{p,l} \left(\frac{u^2}{1+u^2}\right)^l, \\
B_p(u) &= \left(\frac{1}{2m}\right)^p \left(\frac{1}{u}\right)^{p+2} \left(\frac{u^2}{1+u^2}\right)^{p+1} \sum_{l=0}^{p} d_{p,l} \left(\frac{u^2}{1+u^2}\right)^l
\end{aligned} \tag{5.47}
$$

for $k = 0, 1, 2, ...,$ where $c_{k,l}$ and $d_{k,l}$ are constants independent of m and u (but dependent of τ); see [12]. Furthermore, by (5.25), (5.22) and (5.23), it can be proved that

$$
a_p(\tau) = \frac{1}{2\pi i} \int_{C_O \bigcup C_I} h_0(u, \tau) A_p(u) \mathrm{d}u \tag{5.48}
$$

and

$$
b_p(\tau) = \frac{1}{2\pi i} \int_{C_O \bigcup C_I} h_0(u, \tau) B_p(u) \mathrm{d}u; \tag{5.49}
$$

see also [12]. Define $q := \min\{a, b\}$; since in this case both n and $x \to \infty$, and $b^2/a \to 0$, we have $qN \to \infty$, $m/q = O(b/\sqrt{a}) \to 0$ if $a \le b$, and $m/q \sim \arctan\sqrt{a/(1-a)}$ if $a > b$.

Using (5.47), (5.41) and (5.45), it is easily verified that

$$
|A_p(u)| \le \frac{Cm}{b^{p+1}}, \qquad |B_p(u)| \le \frac{C}{b^p} \tag{5.50}
$$

for u on the inner boundary C_I, and

$$
|A_p(u)| \le \frac{Cm}{a^{p+1}}, \qquad |B_p(u)| \le \frac{Cm^2}{a^{p+2}} \tag{5.51}
$$

for u on the outer boundary C_O, where C is used again as a generic symbol for constants independent of u, a and b. Also we have the estimates

$$
\int_{C_I} |\mathrm{d}s| \le \frac{Cm}{b} \quad \text{and} \quad \int_{C_O} |\mathrm{d}s| \le \frac{Ca}{m}. \tag{5.52}
$$

To see this, we take one part of C_I, namely, AB as an example; see (5.35) and (5.36), where Z in (5.35) satisfies $\mathrm{Im}\ Z = \theta b\pi$. Using the second equality in (5.36), and in polar coordinates, $u = re^{i\beta}$, the last equation is equivalent to

$$
\frac{(r^2+1)}{r} \sin\beta = \frac{\theta b\pi}{m}. \tag{5.53}
$$

Thus

$$
\mathrm{d}r = \frac{(1+r^2)^2 m}{(1-r^2)\theta b\pi} \cos\beta \mathrm{d}\beta. \tag{5.54}
$$

Since in this case $r < 1$, we have $|\mathrm{d}r| \le \frac{Cm}{b} |\mathrm{d}\beta|$. By (5.41), $|r| \le C_2 m/b$ and

$$
|\mathrm{d}s| = \sqrt{(\mathrm{d}r)^2 + r^2(\mathrm{d}\beta)^2} \le \frac{Cm}{b} |\mathrm{d}\beta|, \tag{5.55}
$$

from which the first inequality in (5.52) follows. The second inequality can be established in a similar manner.

Since $h_0(u,0) \neq 0$ and $h_0(u,\tau)$ is analytic in τ, we have $|h_0(u,\tau)| \leq C|h_0(u,0)|$ for some constant $C > 0$ and for τ in the neighbourhood of the origin. By a combination of (5.48), (5.49), (5.50), (5.51), (5.52) and (5.46), we obtain

$$|a_p(\tau)| \leq \frac{C}{q^p} \quad \text{and} \quad |b_p(\tau)| \leq \frac{C}{q^p} \tag{5.56}$$

for τ near $\tau = 0$. Furthermore, by (5.24), (5.31), (5.32) and (5.56), we have

$$|c_p| \leq \frac{C}{q^p} \quad \text{and} \quad |d_p| \leq \frac{C}{q^p}. \tag{5.57}$$

To estimate the remainder in (5.33), we split the loop contour into two parts; see Figure 8. The bounded part of the contour, denoted by Γ_1, is contained in a subdomain $\widetilde{D}_u$ of D_u, which has a distance $c_1 a/m$ from the outer boundary of D_u (i.e., C_O), and has a distance $c_2 m/b$ from the inner boundary of D_u (i.e., C_I), c_1 and c_2 being two constants independent of u, a and b. Note that $c_1 a/m$ may become large, whereas $c_2 m/b$ may approach zero. The unbounded part of the contour, denoted by Γ_2, is the rest of the loop outside the subdomain $\widetilde{D}_u$. Put

$$\varepsilon_{p,1}^{+} = \frac{N^{n+1-p}}{2\Gamma(n+1)\pi i} \int_0^{+\infty} \int_{\Gamma_1} \frac{h_p(u,\tau)}{u} e^{Nm\left(u-\frac{1}{u}\right)} \tau^n e^{-N\tau} \mathrm{d}u\mathrm{d}\tau, \tag{5.58}$$

$$\varepsilon_{p,2}^{+} = \frac{N^{n+1-p}}{2\Gamma(n+1)\pi i} \int_0^{+\infty} \int_{\Gamma_2} \frac{h_p(u,\tau)}{u} e^{Nm\left(u-\frac{1}{u}\right)} \tau^n e^{-N\tau} \mathrm{d}u\mathrm{d}\tau. \tag{5.59}$$




Figure 8: The contour in u-plane.

As in [5] and [12], it can be shown that $\varepsilon_{p,2}^{+}$ is exponentially small in comparison with $\varepsilon_{p,1}^{+}$.

To estimate $\varepsilon_{p,1}^{+}$, we also follow [5] and [12] by first deriving an integral representation for $h_k(u,\tau)$. Define recursively

$$\begin{aligned} Q_0(u,w) &= \frac{1}{u-w}, \\ Q_{p+1}(u,w) &= \frac{1}{m}\left(1+\frac{1}{u^2}\right)^{-1}\left\{\frac{1}{u}+\frac{\mathrm{d}}{\mathrm{d}u}\right\}Q_p(u,w) \end{aligned} \tag{5.60}$$

for $p=0,1,2,\cdots$. Using induction, one can easily verify that

$$Q_p(u,w)=\frac{1}{(2m)^p}\sum_{i=0}^{p-1}\sum_{j=0}^{p-i} q_{p,i,j}\left(\frac{1}{1+u^2}\right)^{p+i} u^{2p+i-j}\left(\frac{1}{u-w}\right)^{p+1-i-j}, \tag{5.61}$$

where $q_{p,i,j}$ are some constants depending only on i, j and p. Similar to (4.36), for $w\in\widetilde{D}_u$ we have from (5.22), (5.23) and (5.60) the integral representation

$$h_p(w,\tau)=\frac{1}{2\pi i}\int_{C_I\cup C_O} h_0(u,\tau)Q_p(u,w)\mathrm{d}u. \tag{5.62}$$

From (5.61), it is easy to see that

$$\begin{aligned} |Q_p(u,w)| &\le C\frac{m}{a^{p+1}} \quad \text{for } u\in C_O \text{ and } w\in\widetilde{D}_u, \\ |Q_p(u,w)| &\le C\frac{1}{mb^{p-1}} \quad \text{for } u\in C_I \text{ and } w\in\widetilde{D}_u. \end{aligned} \tag{5.63}$$

Coupling the above inequalities with (5.52) and (5.62), we obtain

$$|h_p(w,\tau)|\le\frac{C}{q^p} \tag{5.64}$$

for $w\in\tilde{D}_u$ and τ in the neighbourhood of $\tau=0$.

If Nm is bounded, one can easily show from (5.58) and (5.64) that

$$|\varepsilon_{p,1}^{+}|=O\left(\frac{1}{q^pN^p}\right). \tag{5.65}$$

If $Nm\to\infty$ we divide the integration path Γ_1 into three pieces: the steepest descent path L_1 through $u=u_+=+i$ in the upper half of the u-plane, the steepest descent path L_2 through $u=u_-=-i$ in the lower half of the u-plane, and the circular arc $|u|=1-\delta_0$, denoted by L_3, which joins the steepest descent paths L_1 and L_2 at $u=u_\delta^+$ and $u=u_\delta^-$, respectively; see Figure 8. Hence,

$$\varepsilon_{p,1}^{+}=I_1+I_2+I_3, \tag{5.66}$$

where I_i, $i=1,2,3$, denotes the integral over the subcontour L_i. Applying the steepest descent method [11, p.84], and using (5.64), it can be easily verified that

$$|I_{1,2}|=O\left(\frac{1}{q^pN^p}\sqrt{\frac{1}{Nm}}\right). \tag{5.67}$$

For I_3, since

$$\begin{aligned}\mathrm{Re}\left(u-\frac{1}{u}\right) &= -\frac{(2-\delta)\delta}{(1-\delta)^2}\mathrm{Re}\ u \le -\frac{(2-\delta)\delta}{(1-\delta)^2}\mathrm{Re}\ u_\delta^{\pm} \\ &= \mathrm{Re}\left(u-\frac{1}{u}\right)\Big|_{u=u_\delta^{\pm}} < \mathrm{Re}\left(u-\frac{1}{u}\right)\Big|_{u=u_{\pm}},\end{aligned}$$

I_2 is exponentially small in comparison with $I_{1,2}$. Therefore,

$$|\varepsilon_{p,1}^{+}| = O\left(\frac{1}{q^p N^p}\sqrt{\frac{1}{Nm}}\right). \tag{5.68}$$

It is well-known that the Bessel function $J_\alpha(t)$ is bounded when t is bounded, and that as $t\to\infty$,

$$J_\alpha(t) \sim \sqrt{\frac{2}{\pi t}}\cos\left(t-\frac{1}{2}\alpha\pi-\frac{1}{4}\pi\right). \tag{5.69}$$

Coupling the estimates (5.65) and (5.68), and together with (5.69) and (5.57), we can find a constant M independent of a and b such that

$$|\varepsilon_p^{+}| \le \frac{M}{qN}\left(|J_0(2Nm)|\frac{|c_{p-1}|}{N^{p-1}} + |J_1(2Nm)|\frac{|d_{p-1}|}{N^{p-1}}\right), \tag{5.70}$$

which establishes the asymptotic nature of the expansion in (5.30), given that $qN\to\infty$. In conclusion, in the case $n^2/xN\to 0$ and $x\to\infty$, (5.30) is an asymptotic expansion.

6 REMAINING CASES

We are now left with only two easy cases to consider.

6.1 $a>0$, a/b^2 large and x bounded

In this case, the series in (1.2) is itself an asymptotic expansion, since

$$\varphi_k(N) := \frac{(-n)_k(-x)_k(n+1)_k}{(-N+1)_k k!k!}, \qquad k=0,1,\cdots,n,$$

is an asymptotic sequence when $xN/n^2\to\infty$; see Table 1 and also [11, p.10].

6.2 $a<0$

The case $a<0$ is relatively easy, compared with the case $a>0$. Let us start with the integral representation (2.23). For fixed $w\in\gamma_2$, the phase function $\widetilde{f}(t,w)$ in (2.23) has a saddle point

$$t_0(w) = \frac{2w-1+\sqrt{1+4b^2(w-1)w}}{2(1+b)w}; \tag{6.1}$$

cf.(2.5). Note that the only difference between $f(t,w)$ in (2.2) and $\widetilde{f}(t,w)$ in (2.24) is the choice of branch cuts in the w-plane due to the term $a\ln w - a\ln(w-1)$ in $f(t,w)$ and the term $a\ln(-w) - a\ln(1-w)$ in $\widetilde{f}(t,w)$. Similar to (3.1), for any fixed $w \in \gamma_2$, we have $t_0(w) = 1 - b + O(b^2)$ as $b \to 0$; in particular, $t_0(0) = 1 - b$. By the same reasoning given for (3.2), we introduce the mapping $t \to \tau$ defined by

$$b\ln\tau - \tau + A = \widetilde{f}(t,w) - \widetilde{f}(t_0(w),w) \tag{6.2}$$

with

$$\tau(t_0(w)) = b. \tag{6.3}$$

Coupling (6.3) and (6.2) yields $A = b - b\ln b$. From (6.2), we have

$$\begin{aligned}
&\frac{\mathrm{d}t}{\mathrm{d}\tau} = \frac{b/\tau - 1}{\partial\widetilde{f}(t,w)/\partial t} = \frac{(\tau-b)t(1-t)[1-(1-t)w]}{\tau(1+b)w(t-t_0^+(w))(t-t_0^-(w))}, && \tau \neq b,\\
&\frac{\mathrm{d}t}{\mathrm{d}\tau} = \left\{\frac{t_0^+(w)(1-t_0^+(w))[1-(1-t_0^+(w))w]}{b\sqrt{1+4b^2(w-1)w}}\right\}^{1/2}, && \tau = b,
\end{aligned} \tag{6.4}$$

where we have used L'Hôspital's rule for $\tau = b$ and

$$t_0^{\pm}(w) = \frac{2w - 1 \pm \sqrt{1+4b^2(w-1)w}}{2(1+b)w}; \tag{6.5}$$

cf.(3.5). Furthermore, it follows from (6.2) and (2.23) that

$$\begin{aligned}
t_n(x,N+1) =& \frac{(-1)^n}{2\pi i}\frac{\Gamma(n+N+2)e^n b^{-n}}{\Gamma(n+1)\Gamma(N-n+1)}\\
&\times \int_0^{+\infty}\int_{\gamma_2} \frac{1}{w-1} e^{N\widetilde{f}(t_0(w),w)} e^{N(b\ln\tau - \tau)} \frac{\mathrm{d}t}{\mathrm{d}\tau}\mathrm{d}w\mathrm{d}\tau;
\end{aligned} \tag{6.6}$$

cf.(3.6). Solving the equation $\partial f(t_0(w),w)/\partial w = 0$, we obtain the saddle points

$$\begin{aligned}
w_{\pm} =& \frac{b \pm \sqrt{b^2 - 4a + 4a^2}}{2b}\\
=& \frac{1}{2} \pm \frac{1}{2}\sqrt{1 - \frac{4a(1-a)}{b^2}};
\end{aligned} \tag{6.7}$$

cf.(2.9). The relevant saddle point on the integral path γ_2 is the negative saddle point w_-.

The following procedure is the same as that given in [8]. Recall the Hankel integral for the Gamma function

$$\frac{1}{\Gamma(z)} = \frac{1}{2\pi i}\int_{-\infty}^{(0+)} e^u u^{-z}\mathrm{d}u, \tag{6.8}$$

where the contour is a loop starting at $-\infty$, encircling the origin in the counterclockwise direction and returning to $-\infty$. With $z = -aN + 1$ and u replaced by $-u$, we obtain

$$\frac{1}{\Gamma(-aN+1)} = \frac{N^{aN}}{2\pi i}\int_{\infty}^{(0+)} \frac{1}{u} e^{N\widetilde{\psi}(u)}\mathrm{d}u, \tag{6.9}$$

where $\widetilde{\psi}(u) = a\ln(-u) - u$. Make the transformation $w \to u$ defined by

$$\widetilde{f}(t_0(w), w) = a\ln(-u) - u + \gamma \tag{6.10}$$

with

$$u(w_-) = a, \tag{6.11}$$

where γ is a constant to be determined. We have from (6.10)

$$\begin{aligned} \frac{\mathrm{d}w}{\mathrm{d}u} &= \frac{(a-u)w(w-1)\left[(1-2a) + \sqrt{4b^2w^2 - 4b^2w + 1}\right]}{2ub^2(w-w_+)(w-w_-)}, \qquad u \neq a, \\ \frac{\mathrm{d}w}{\mathrm{d}u} &= \left\{\frac{(1-a)(1-2a)}{b^3\sqrt{b^2-4a+4a^2}}\right\}^{1/2}, \qquad u = a, \end{aligned} \tag{6.12}$$

where we again have used L'H$\hat{o}$spital's rule for $u = a$.

By (6.6) and (6.10), we have

$$\begin{aligned} t_n(x, N+1) =& \frac{(-1)^n}{2\pi i}\frac{\Gamma(n+N+2)e^n b^{-n}e^{N\gamma}}{\Gamma(n+1)\Gamma(N-n+1)} \\ & \times \int_0^{+\infty}\int_\infty^{(0^+)} \frac{h(u,\tau)}{u} e^{N(a\ln(-u)-u)} e^{N(b\ln\tau - \tau)}\mathrm{d}u\mathrm{d}\tau, \end{aligned} \tag{6.13}$$

where

$$h(u,\tau) = \frac{u}{w-1}\frac{\mathrm{d}t}{\mathrm{d}\tau}\frac{\mathrm{d}w}{\mathrm{d}u}. \tag{6.14}$$

Define recursively

$$h_l(u,\tau) = a_l(\tau) + (u-a)g_l(u,\tau), \tag{6.15}$$

$$h_{l+1}(u,\tau) = u\frac{\partial g_l(u,\tau)}{\partial u}, \tag{6.16}$$

where $h_0(u,\tau) = h(u,\tau)$, and expand $a_l(\tau)$ into a Maclaurin series

$$a_l(\tau) = \sum_{j=0}^{\infty} a_{l,j}\tau^j. \tag{6.17}$$

By an integration-by-parts procedure, we obtain

$$t_n(x, N+1) = \frac{(-1)^n\Gamma(n+N+2)e^n b^{-n}e^{N\gamma}}{\Gamma(N-n+1)\Gamma(-aN+1)N^{aN+n+1}}\left[\sum_{l=0}^{p-1}\frac{c_l}{N^l} + \varepsilon_p^-\right], \tag{6.18}$$

where

$$c_l = \frac{N^{n+1}}{\Gamma(n+1)}\int_0^\infty a_l(\tau)\tau^n e^{-N\tau}\mathrm{d}\tau \sim \sum_{m=0}^{\infty} a_{l,m}\frac{\Gamma(n+m+1)}{\Gamma(n+1)N^m} \tag{6.19}$$

and

$$\varepsilon_p^- = \frac{N^{aN+n+1-p}\Gamma(-aN+1)}{2\Gamma(n+1)\pi i}\int_0^{+\infty}\int_{-\infty}^{(0^+)} \frac{h_p(u,\tau)}{u} e^{N\widetilde{\psi}(u)}\tau^n e^{-N\tau}\mathrm{d}u\mathrm{d}\tau. \tag{6.20}$$

Estimation of the error term given in (6.20) is exactly the same as what we have done in [8].

References

[1] R. Beals and R. Wong, *Special Functions, A Graduate Text*, Cambridge University Press, Cambridge, 2010.

[2] C. Chester, B. Friedman and F. Ursell, *An extension of the method of steepest descents*, Proc. Cambridge Philos. Soc. **53** (1957), 599-611.

[3] X.-S. Jin and R. Wong, *Uniform asymptotic expansions for Meixner polynomials*, Constr. Approx. **14** (1998), 113-150.

[4] Y. Lin and R. Wong, *Global Asymptotics of the Hahn Polynomials*, Analysis and Applications, to appear.

[5] A. B. Olde Daalhuis and N. M. Temme, *Uniform Airy-type expansions of integrals*, SIAM J. Math. Anal. **25** (1994), 304-321.

[6] F. W. J. Olver, *Asymptotics and Special Functions*, Academic Press, New York, 1974. (Reprinted by A. K. Peters, Wellesley, MA, 1997.)

[7] F. W. J. Olver, D. W. Lozier, C. W. Clark and R. F. Boisvert, *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, 2010.

[8] J. H. Pan and R. Wong, *Uniform Asymptotic Expansions for the Discrete Chebyshev Polynomials*, Studies in Applied Mathematics **128** (2012), 337-384.

[9] Bo, Rui and R. Wong, *Uniform asymptotic expansion of Charlier polynomials*. Methods and Applications of Analysis **1** (1994), 294-313.

[10] I. I. Sharapudinov, *Asymptotic Properties of Orthogonal Hahn Polynomials in a Discrete Variable*, Sbornik: Mathematics, Vol. 68, **1** (1991), 111-132.

[11] R. Wong, *Asymptotic Approximations of Integrals*, Academic Press, Boston, 1989. (Reprinted by SIAM, Philadelphia, PA, 2001)

[12] R. Wong and Y. Q. Zhao, *Estimates for the error term in a uniform asymptotic expansion of the Jacobi polynomials*. Anal. Appl. (Singap.) **1** (2003), no. 2, 213-241.

[13] R. Wong, *Lecture Notes on Applied Analysis*, World Scientific, Singapore, 2010.